

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CDX2 (GI Epithelial Marker) (CDX2/2951R), CF405S conjugate, 0.1mg/mL, Clone: [CDX2/2951R], Rabbit, Monoclonal BOT-BNC042951-100**

Artikelname	CDX2 (GI Epithelial Marker) (CDX2/2951R), CF405S conjugate, 0.1mg/mL, Clone: [CDX2/2951R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042951-100
Hersteller Artikelnummer	BNC042951-100
Alternativnummer	BOT-BNC042951-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa150-249) of human CDX2 protein (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CD...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[CDX2/2951R]
Molekulargewicht	40 kDa
UniProt	<a href="#">Q99626</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user