

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

von Willebrand Factor(VWF635), CF640R conjugate, 0.1mg/mL, Clone: [VWF635], Mouse, Monoclonal BOT-BNC400635-500

Artikelname	von Willebrand Factor(VWF635), CF640R conjugate, 0.1mg/mL, Clone: [VWF635], Mouse, Monoclonal
Artikelnummer	BOT-BNC400635-500
Hersteller Artikelnummer	BNC400635-500
Alternativnummer	BOT-BNC400635-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human vWF fragment
Konjugation	CF640R
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[VWF635]

Molekulargewicht	250 kDa
UniProt	P04275
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user