

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Pmel17 / gp100 / SILV(PMEL/783), CF640R conjugate, 0.1mg/mL, Clone: [PMEL/783], Mouse, Monoclonal BOT-BNC400783-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Pmel17 / gp100 / SILV(PMEL/783), CF640R conjugate, 0.1mg/mL, Clone: [PMEL/783], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC400783-500   |
| Hersteller Artikelnummer | BNC400783-500   |
| Alternativnummer         | BOT-BNC400783-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human SILV protein  |
| Konjugation              | CF640R  |
| Produktbeschreibung      | Cytotoxic T lymphocytes (CTLs) recognize melanoma-associated antigens, which belong to three main groups. These groups include tumor-associated testis-specific antigens, melanocyte differentiation antigens and mutated or aberrantly expressed antigens... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [PMEL/783]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 90-100 kDa   |
| UniProt                | <a href="#">P40967</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |