

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgA Secretory Component(ECM1/792), CF640R conjugate, 0.1mg/mL, Clone: [ECM1/792], Mouse, Monoclonal BOT-BNC400792-500

| | |
|--------------------------|---|
| Artikelname | IgA Secretory Component(ECM1/792), CF640R conjugate, 0.1mg/mL, Clone: [ECM1/792], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC400792-500 |
| Hersteller Artikelnummer | BNC400792-500 |
| Alternativnummer | BOT-BNC400792-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human ECM1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ECM1/792] |

| | |
|------------------------|--|
| Molekulargewicht | ~80 kDa |
| UniProt | Q16610 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |