

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD56 / NCAM(NCAM1/795), CF640R conjugate, 0.1mg/mL, Clone: [NCAM1/795], Mouse, Monoclonal BOT-BNC400795-100

Artikelname	CD56 / NCAM(NCAM1/795), CF640R conjugate, 0.1mg/mL, Clone: [NCAM1/795], Mouse, Monoclonal
Artikelnummer	BOT-BNC400795-100
Hersteller Artikelnummer	BNC400795-100
Alternativnummer	BOT-BNC400795-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human NCAM1 protein
Konjugation	CF640R
Produktbeschreibung	This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NCAM1/795]
Molekulargewicht	180, 145 and 125 kDa
UniProt	P13591
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p>