

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), CF640R conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal BOT-BNC400854-500

| | |
|--------------------------|---|
| Artikelname | CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), CF640R conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC400854-500 |
| Hersteller Artikelnummer | BNC400854-500 |
| Alternativnummer | BOT-BNC400854-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human, Monkey, Primate |
| Immunogen | Human tonsil B cells (L26), Recombinant human MS4A1 protein (IGEL/773) |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [L26 IGEL/773] |

| | |
|------------------------|---|
| Molekulargewicht | 33-37 kDa |
| UniProt | P11836 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Does not react with Cow, Dog, Pig or Rat For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |