

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Glucose Regulated Protein 94 (GRP94)(HSP90B1/1192), CF640R conjugate, 0.1mg/mL, Clone: [HSP90B1/1192], Rat, Monoclonal BOT-BNC401192-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Glucose Regulated Protein 94 (GRP94)(HSP90B1/1192), CF640R conjugate, 0.1mg/mL, Clone: [HSP90B1/1192], Rat, Monoclonal  |
| Artikelnummer            | BOT-BNC401192-500   |
| Hersteller Artikelnummer | BNC401192-500   |
| Alternativnummer         | BOT-BNC401192-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rat   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human HSP90B1 protein   |
| Konjugation              | CF640R  |
| Produktbeschreibung      | Recognizes a protein of 94 kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This MAb is highly specifi... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [HSP90B1/1192]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 94 kDa  |
| UniProt                | <a href="#">P14625</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 0.5-1.0 ug/mL Western blotting 0.5-1.0 ug/mL Predicted to show broad species reactivity Optimal dilution for a specific application should be determined by user |