

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HPV-16 (Human Papilloma Virus 16) (HPV16/1296), CF640R conjugate, 0.1mg/mL, Clone: [HPV16/1296], Mouse, Monoclonal BOT-BNC401296-500

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | HPV-16 (Human Papilloma Virus 16) (HPV16/1296), CF640R conjugate, 0.1mg/mL, Clone: [HPV16/1296], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401296-500 |
| Hersteller Artikelnummer | BNC401296-500 |
| Alternativnummer | BOT-BNC401296-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Virus |
| Immunogen | Human papilloma virus type 16 |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody reacts with E6 protein of human papilloma virus type 16 (HPV-16). Infection with specific types of HPV has been associated with an increased risk of developing cervical neoplasia. HPV types 6 and 11 have been associated with relatively ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [HPV16/1296] |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht | kDa |
| UniProt | Not Applicable |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |