

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MAML2 (Mastermind Like Transcriptional Coactivator 2) (MAML2/1302), CF640R conjugate, 0.1mg/mL, IgG2a, Clone: [MAML2/1302], Mouse, Monoclonal BOT-BNC401302-500

| | |
|--------------------------|---|
| Artikelname | MAML2 (Mastermind Like Transcriptional Coactivator 2) (MAML2/1302), CF640R conjugate, 0.1mg/mL, IgG2a, Clone: [MAML2/1302], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401302-500 |
| Hersteller Artikelnummer | BNC401302-500 |
| Alternativnummer | BOT-BNC401302-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length MAML2 |
| Konjugation | CF640R |
| Produktbeschreibung | MAML2 (mastermind-like protein 2), also known as MAM2, MAM3 or MLLMAML2, is a nuclear speckle protein that acts as a transcriptional co-activator for Notch receptors. The Notch signaling pathway influences cell fate by regulating the ability of precu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [MAML2/1302] |
| Molekulargewicht | 125 kDa |
| Isotyp | IgG2a |
| UniProt | Q8IZL2 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |