

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **VLDL-Receptor (Very Low Density Lipoprotein Receptor) (VLDLR/1337), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [VLDLR/1337], Mouse, Monoclonal BOT-BNC401337-500**

Artikelname	VLDL-Receptor (Very Low Density Lipoprotein Receptor) (VLDLR/1337), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [VLDLR/1337], Mouse, Monoclonal
Artikelnummer	BOT-BNC401337-500
Hersteller Artikelnummer	BNC401337-500
Alternativnummer	BOT-BNC401337-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human VLDLR fragment from c-terminal (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	VLDLR (very low density lipoprotein receptor) is a member of the LDL receptor gene family, which includes LDL receptor, LRP, megalin, VLDLR and ApoER2. The LDL receptor family is characterized by a cluster of cysteine-rich class A repeats, epidermal ...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[VLDLR/1337]
Molekulargewicht	143-161 kDa
Isotyp	IgG1
UniProt	<a href="#">P98155</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user