

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8/18 (KRT8.18/1346), CF640R conjugate, 0.1mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal BOT-BNC401346-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 8/18 (KRT8.18/1346), CF640R conjugate, 0.1mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC401346-500   |
| Hersteller Artikelnummer | BNC401346-500   |
| Alternativnummer         | BOT-BNC401346-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Cytoskeleton preparation from HeLa cells  |
| Konjugation              | CF640R  |
| Produktbeschreibung      | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb recognizes all simple epithelia including glandular epithelium, for example thyr... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [KRT8.18/1346]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 52.5 kDa (CK8), 45 kDa (CK18)   |
| UniProt                | <a href="#">P05787</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |