

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Luteinizing Hormone / Choriogonadotropin Receptor (LHCGR) (LHCGR/1416), CF640R conjugate, 0.1mg/mL, Clone: [LHCGR/1416], Mouse, Monoclonal BOT-BNC401416-500

| | |
|--------------------------|---|
| Artikelname | Luteinizing Hormone / Choriogonadotropin Receptor (LHCGR) (LHCGR/1416), CF640R conjugate, 0.1mg/mL, Clone: [LHCGR/1416], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401416-500 |
| Hersteller Artikelnummer | BNC401416-500 |
| Alternativnummer | BOT-BNC401416-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Purified recombinant N-terminal fragment of human LHCGR (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a protein of 78-85 kDa, which is identified as luteinizing hormone/choriogonadotropin receptor. Luteinizing hormone plays a role in spermatogenesis and ovulation by stimulating the testes and ovaries to produce steroids. Chor... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [LHCGR/1416] |
| Molekulargewicht | 78-85 kDa |
| UniProt | P22888 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |