

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MCM7 (Proliferation Marker) (MCM7/1467), CF640R conjugate, 0.1mg/mL, Clone: [MCM7/1467], Mouse, Monoclonal BOT-BNC401467-100

| | |
|--------------------------|---|
| Artikelname | MCM7 (Proliferation Marker) (MCM7/1467), CF640R conjugate, 0.1mg/mL, Clone: [MCM7/1467], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401467-100 |
| Hersteller Artikelnummer | BNC401467-100 |
| Alternativnummer | BOT-BNC401467-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human MCM7 protein fragment (aa195-319) (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication c... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [MCM7/1467] |
| Molekulargewicht | 88 kDa |
| UniProt | P33993 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |