

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MCM7 (Proliferation Marker) (MCM7/1468), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [MCM7/1468], Mouse, Monoclonal BOT-BNC401468-500**

Artikelname	MCM7 (Proliferation Marker) (MCM7/1468), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [MCM7/1468], Mouse, Monoclonal
Artikelnummer	BOT-BNC401468-500
Hersteller Artikelnummer	BNC401468-500
Alternativnummer	BOT-BNC401468-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human MCM7 protein fragment (aa195-319) (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication c...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[MCM7/1468]
Molekulargewicht	88 kDa
Isotyp	IgG2b
UniProt	<a href="#">P33993</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user