

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/1487R), CF640R conjugate, 0.1mg/mL, Clone: [DG1/1487R], Rabbit, Monoclonal BOT-BNC401487-100**

Artikelname	DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/1487R), CF640R conjugate, 0.1mg/mL, Clone: [DG1/1487R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC401487-100
Hersteller Artikelnummer	BNC401487-100
Alternativnummer	BOT-BNC401487-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human DOG-1 protein
Konjugation	CF640R
Produktbeschreibung	Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST s), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to d...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[DG1/1487R]

Molekulargewicht	~114 kDa
UniProt	<a href="#">Q5XXA6</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry