

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Catenin, beta (CTNNB1) (CTNNB1/2030R), CF640R conjugate, 0.1mg/mL, Clone: [CTNNB1/2030R], Rabbit, Monoclonal BOT-BNC402030-100

| | |
|--------------------------|---|
| Artikelname | Catenin, beta (CTNNB1) (CTNNB1/2030R), CF640R conjugate, 0.1mg/mL, Clone: [CTNNB1/2030R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC402030-100 |
| Hersteller Artikelnummer | BNC402030-100 |
| Alternativnummer | BOT-BNC402030-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human i-catenin protein |
| Konjugation | CF640R |
| Produktbeschreibung | Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CTNNB1/2030R] |

| | |
|------------------------|--|
| Molekulargewicht | 92 kDa |
| UniProt | P35222 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user |