

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), CF640R conjugate, 0.1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal BOT-BNC402034-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), CF640R conjugate, 0.1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal   |
| Artikelnummer            | BOT-BNC402034-100   |
| Hersteller Artikelnummer | BNC402034-100   |
| Alternativnummer         | BOT-BNC402034-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human CD22 protein (exact sequence is proprietary)  |
| Konjugation              | CF640R  |
| Produktbeschreibung      | This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [LPFS2/2034R]   |
| Molekulargewicht       | 120 kDa   |
| UniProt                | <a href="#">P26842</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |