

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) (PMEL/2037), CF640R conjugate, 0.1mg/mL, Clone: [PMEL/2037], Mouse, Monoclonal BOT-BNC402037-100

| | |
|--------------------------|---|
| Artikelname | gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) (PMEL/2037), CF640R conjugate, 0.1mg/mL, Clone: [PMEL/2037], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402037-100 |
| Hersteller Artikelnummer | BNC402037-100 |
| Alternativnummer | BOT-BNC402037-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human SILV protein fragment (around aa 376-502) (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | Cytotoxic T lymphocytes (CTLs) recognize melanoma-associated antigens, which belong to three main groups. These groups include tumor-associated testis-specific antigens, melanocyte differentiation antigens and mutated or aberrantly expressed antigens... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [PMEL/2037] |
| Molekulargewicht | 90-100 kDa |
| UniProt | P40967 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |