

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) (KRT16/2043R), CF640R conjugate, 0.1mg/mL, Clone: [KRT16/2043R], Rabbit, Monoclonal BOT-BNC402043-100**

Artikelname	Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) (KRT16/2043R), CF640R conjugate, 0.1mg/mL, Clone: [KRT16/2043R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC402043-100
Hersteller Artikelnummer	BNC402043-100
Alternativnummer	BOT-BNC402043-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment from the C-terminal of human Cytokeratin 16
Konjugation	CF640R
Produktbeschreibung	Cytokeratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I Cytokeratins (Cytokeratins 9 to 23) and one type II Cytokeratins (keratins 1 to 8). The cytokeratin proteins play a cr...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[KRT16/2043R]
Molekulargewicht	48 kDa
UniProt	<a href="#">P08779</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user