

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cyclin E (G1/S-Phase Cyclin) (CCNE1/2460), CF640R conjugate, 0.1mg/mL, Clone: [CCNE1/2460], Mouse, Monoclonal BOT-BNC402460-100**

Artikelname	Cyclin E (G1/S-Phase Cyclin) (CCNE1/2460), CF640R conjugate, 0.1mg/mL, Clone: [CCNE1/2460], Mouse, Monoclonal
Artikelnummer	BOT-BNC402460-100
Hersteller Artikelnummer	BNC402460-100
Alternativnummer	BOT-BNC402460-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Cyclin E (CCNE1) protein fragment (around aa 10-176) (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	Cyclin E belongs to the highly conserved cyclin family, whose members exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. Cyclins function as regulators of CDK kinases. Cyclin E fo...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[CCNE1/2460]
Molekulargewicht	53 kDa
UniProt	<a href="#">P24864</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user