

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) (NGFR/2550R), CF640R conjugate, 0.1mg/mL, Clone: [NGFR/2550R], Rabbit, Monoclonal BOT-BNC402550-500**

Artikelname	NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) (NGFR/2550R), CF640R conjugate, 0.1mg/mL, Clone: [NGFR/2550R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC402550-500
Hersteller Artikelnummer	BNC402550-500
Alternativnummer	BOT-BNC402550-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human p75 NGFR protein fragment (around aa 281-421) (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	This antibody recognizes a glycoprotein of 75 kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melano...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NGFR/2550R]
Molekulargewicht	75 kDa
UniProt	<a href="#">P08138</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user