

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AKT1 (Prognostic Marker for Neuroendocrine Tumors) (AKT1/2552), CF640R conjugate, 0.1mg/mL, Clone: [AKT1/2552], Mouse, Monoclonal BOT-BNC402552-100

| | |
|--------------------------|---|
| Artikelname | AKT1 (Prognostic Marker for Neuroendocrine Tumors) (AKT1/2552), CF640R conjugate, 0.1mg/mL, Clone: [AKT1/2552], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402552-100 |
| Hersteller Artikelnummer | BNC402552-100 |
| Alternativnummer | BOT-BNC402552-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human AKT1 protein (around aa 85-189) (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a protein of 62 kDa, which is identified as AKT1. The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKBbeta or RacPK-beta) and Akt3 (also des... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [AKT1/2552] |
| Molekulargewicht | 62 kDa |
| UniProt | P31749 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |