

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NKX6.1 (Marker for Pancreatic and Duodenal Neuroendocrine Tumors) (NKX61/2561), CF640R conjugate, 0.1mg/mL, Clone: [NKX61/2561], Mouse, Monoclonal BOT-BNC402561-100

| | |
|--------------------------|---|
| Artikelname | NKX6.1 (Marker for Pancreatic and Duodenal Neuroendocrine Tumors) (NKX61/2561), CF640R conjugate, 0.1mg/mL, Clone: [NKX61/2561], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402561-100 |
| Hersteller Artikelnummer | BNC402561-100 |
| Alternativnummer | BOT-BNC402561-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human full-length recombinant NKX6.1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | Members of the Nkx family of homeodomain proteins are key regulators of growth and development in several tissues, including brain, heart and pancreas. During neural development, sonic hedgehog (Shh) is known to control cell fate and mitogenesis, whi... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [NKX61/2561] |
| Molekulargewicht | 40-50 kDa |
| UniProt | P78426 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |