

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Desmoglein-3 (Squamous Cell Marker)(DSG3/2796), CF640R conjugate, 0.1mg/mL, Clone: [DSG3/2796], Mouse, Monoclonal BOT-BNC402796-500**

Artikelname	Desmoglein-3 (Squamous Cell Marker)(DSG3/2796), CF640R conjugate, 0.1mg/mL, Clone: [DSG3/2796], Mouse, Monoclonal
Artikelnummer	BOT-BNC402796-500
Hersteller Artikelnummer	BNC402796-500
Alternativnummer	BOT-BNC402796-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 379-491) human DSG3 protein (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	Recognizes a protein of 130 kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of de...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[DSG3/2796]

Molekulargewicht	130 kDa
UniProt	<a href="#">P32926</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry