

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Phosphotyrosine (P-Tyr)(PY2870R), CF640R conjugate, 0.1mg/mL, Clone: [PY2870R], Rabbit, Monoclonal BOT-BNC402870-500

| | |
|--------------------------|---|
| Artikelname | Phosphotyrosine (P-Tyr)(PY2870R), CF640R conjugate, 0.1mg/mL, Clone: [PY2870R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC402870-500 |
| Hersteller Artikelnummer | BNC402870-500 |
| Alternativnummer | BOT-BNC402870-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Spezies Reaktivität | All |
| Immunogen | Recombinant full-length human phosphotyrosine protein |
| Konjugation | CF640R |
| Produktbeschreibung | Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [PY2870R] |
| Molekulargewicht | Depends upon the phosphorylated target |

| | |
|------------------------|--|
| UniProt | Not Applicable |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |