

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326(VU-1D9), CF647 conjugate, 0.1mg/mL, Clone: [VU-1D9], Mouse, Monoclonal BOT-BNC470017-100

Artikelname	Ep-CAM / CD326(VU-1D9), CF647 conjugate, 0.1mg/mL, Clone: [VU-1D9], Mouse, Monoclonal
Artikelnummer	BOT-BNC470017-100
Hersteller Artikelnummer	BNC470017-100
Alternativnummer	BOT-BNC470017-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	Exosome Staining, FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Small cell lung carcinoma cells
Konjugation	CF647
Produktbeschreibung	This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[VU-1D9]

Molekulargewicht	40-43 kDa
UniProt	P16422
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat or ferret, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user