

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Kappa Light Chain(L1C1), CF647 conjugate, 0.1mg/mL, Clone: [L1C1], Mouse, Monoclonal BOT-BNC470018-500

| | |
|--------------------------|---|
| Artikelname | Human Kappa Light Chain(L1C1), CF647 conjugate, 0.1mg/mL, Clone: [L1C1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470018-500 |
| Hersteller Artikelnummer | BNC470018-500 |
| Alternativnummer | BOT-BNC470018-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Human B-Lymphoma Cells |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [L1C1] |

| | |
|------------------------|--|
| Molekulargewicht | ~22.5 kDa |
| UniProt | P01601 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-2.0 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user |