

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p21 / WAF1(WA-1), CF647 conjugate, 0.1mg/mL, Clone: [WA-1], Mouse, Monoclonal BOT-BNC470043-500**

|                          |   |
|--------------------------|---|
| Artikelname              | p21 / WAF1(WA-1), CF647 conjugate, 0.1mg/mL, Clone: [WA-1], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC470043-500   |
| Hersteller Artikelnummer | BNC470043-500   |
| Alternativnummer         | BOT-BNC470043-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human, Monkey, Mouse, Primate, Rat  |
| Immunogen                | Human recombinant p21 protein   |
| Konjugation              | CF647   |
| Produktbeschreibung      | This MAb recognizes a 21 kDa protein, identified as the p21WAF1 tumor suppressor protein. This MAb is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdk s and a tum... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [WA-1]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 21 kDa   |
| UniProt                | <a href="#">P38936</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |