

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HLA-DRB(LN-3), CF647 conjugate, 0.1mg/mL, Clone: [LN-3], Mouse, Monoclonal BOT-BNC470057-500

| | |
|--------------------------|---|
| Artikelname | HLA-DRB(LN-3), CF647 conjugate, 0.1mg/mL, Clone: [LN-3], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470057-500 |
| Hersteller Artikelnummer | BNC470057-500 |
| Alternativnummer | BOT-BNC470057-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Monkey |
| Immunogen | Activated human peripheral blood mononuclear cells |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb reacts with a 28 kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do no... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [LN-3] |

| | |
|------------------------|---|
| Molekulargewicht | 28 kDa (beta chain) |
| UniProt | P01911 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Does not react with mouse, others not known Optimal dilution for a specific application should be determined by user</p> |