

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Histone H1(AE-4), CF647 conjugate, 0.1mg/mL, Clone: [AE-4], Mouse, Monoclonal BOT-BNC470096-500

| | |
|--------------------------|---|
| Artikelname | Histone H1(AE-4), CF647 conjugate, 0.1mg/mL, Clone: [AE-4], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470096-500 |
| Hersteller Artikelnummer | BNC470096-500 |
| Alternativnummer | BOT-BNC470096-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ChIP, FC, IF, IHC, IP, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Nuclei of human leukemia biopsy cells |
| Konjugation | CF647 |
| Produktbeschreibung | Eukaryotic histones are basic, water-soluble nuclear proteins that form hetero-octameric nucleosome particles. They wrap 146 base pairs of DNA in a left-handed supealpha-helicalturn sequentially to form chromosomal fiber. Two molecules of each of the... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [AE-4] |

| | |
|------------------------|--|
| Molekulargewicht | ~30 kDa |
| UniProt | Multiple |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |