

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(GP1.4), CF647 conjugate, 0.1mg/mL, Clone: [GP1.4], Mouse, Monoclonal BOT-BNC470159-500**

Artikelname	Mucin 1 / EMA / Episialin / CD227(GP1.4), CF647 conjugate, 0.1mg/mL, Clone: [GP1.4], Mouse, Monoclonal
Artikelnummer	BOT-BNC470159-500
Hersteller Artikelnummer	BNC470159-500
Alternativnummer	BOT-BNC470159-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Human milk fat globule membranes
Konjugation	CF647
Produktbeschreibung	In Western blotting, this antibody recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. This MAAb reacts with the DTRP epitope in the tandem repeats. The alpha subunit has cell adhesive properties. It can act both ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[GP1.4]

Molekulargewicht	265-400 kDa
UniProt	<a href="#">P15941</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user