

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

GFAP(GA-5), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [GA-5], Mouse, Monoclonal BOT-BNC470167-500

Artikelname	GFAP(GA-5), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [GA-5], Mouse, Monoclonal
Artikelnummer	BOT-BNC470167-500
Hersteller Artikelnummer	BNC470167-500
Alternativnummer	BOT-BNC470167-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Bovine, Gallus, Human, Mouse, Porcine, Rabbit, Rat
Immunogen	GFAP isolated from pig spinal cord
Konjugation	CF647
Produktbeschreibung	This MAb recognizes a protein of ~50 kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for I...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[GA-5]

Molekulargewicht	~50 kDa
Isotyp	IgG1
UniProt	P14136
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p>