

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD50 (ICAM-3)(101-1D2), CF647 conjugate, 0.1mg/mL, Clone: [101-1D2], Mouse, Monoclonal BOT-BNC470177-100

| | |
|--------------------------|---|
| Artikelname | CD50 (ICAM-3)(101-1D2), CF647 conjugate, 0.1mg/mL, Clone: [101-1D2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470177-100 |
| Hersteller Artikelnummer | BNC470177-100 |
| Alternativnummer | BOT-BNC470177-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, Functional Studies |
| Spezies Reaktivität | Human |
| Immunogen | Stimulated human leukocytes |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes an N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (WS: IV & V). Its epitope localizes in the D2 extracellular domain and is resistant to neuraminidase and proteases. CD50 is the major ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [101-1D2] |

| | |
|------------------------|---|
| Molekulargewicht | 110-160 kDa |
| UniProt | P32942 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin/paraffin tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |