

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD50 (ICAM-3)(186-2G9), CF647 conjugate, 0.1mg/mL, Clone: [186-2G9], Mouse, Monoclonal BOT-BNC470178-100

| | |
|--------------------------|---|
| Artikelname | CD50 (ICAM-3)(186-2G9), CF647 conjugate, 0.1mg/mL, Clone: [186-2G9], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470178-100 |
| Hersteller Artikelnummer | BNC470178-100 |
| Alternativnummer | BOT-BNC470178-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Stimulated human leukocytes |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes the D1 domain of an N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (Leucocyte Workshop VI). CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhe... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [186-2G9] |

| | |
|------------------------|---|
| Molekulargewicht | 110-160 kDa |
| UniProt | P32942 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Blocks binding of CD11a LFA-1 to CD50 ICAM-3 Optimal dilution for a specific application should be determined by user |