

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27 / KIP1(SX53G8), CF647 conjugate, 0.1mg/mL, Clone: [SX53G8], Mouse, Monoclonal BOT-BNC470669-500

Artikelname	p27 / KIP1(SX53G8), CF647 conjugate, 0.1mg/mL, Clone: [SX53G8], Mouse, Monoclonal
Artikelnummer	BOT-BNC470669-500
Hersteller Artikelnummer	BNC470669-500
Alternativnummer	BOT-BNC470669-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, IP, WB
Spezies Reaktivität	Human, Monkey, Mouse, Rat
Immunogen	Purified GST-p27 fusion protein of human origin
Konjugation	CF647
Produktbeschreibung	This MAb recognizes a 27 kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human brea...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[SX53G8]

Molekulargewicht	25-26 kDa
UniProt	P46527
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user