

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8(H1 + TS1), CF647 conjugate, 0.1mg/mL, Clone: [H1 TS1], Mouse, Monoclonal BOT-BNC470687-500

Artikelname	Cytokeratin 8(H1 + TS1), CF647 conjugate, 0.1mg/mL, Clone: [H1 TS1], Mouse, Monoclonal
Artikelnummer	BOT-BNC470687-500
Hersteller Artikelnummer	BNC470687-500
Alternativnummer	BOT-BNC470687-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Cytoskeleton preparation containing cytokeratin 8 (H1), Keratin preparation from a human carcinoma (TS1)
Konjugation	CF647
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[H1 TS1]
Molekulargewicht	52.5 kDa
UniProt	P05787
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user