

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Chromogranin A(LK2H10 + PHE5), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [LK2H10 PHE5], Mouse, Monoclonal BOT-BNC470698-100

Artikelname	Chromogranin A(LK2H10 + PHE5), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [LK2H10 PHE5], Mouse, Monoclonal
Artikelnummer	BOT-BNC470698-100
Hersteller Artikelnummer	BNC470698-100
Alternativnummer	BOT-BNC470698-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Monkey, Mouse, Porcine, Rat
Immunogen	Human pheochromocytoma (LK2H10 & PHE5)
Konjugation	CF647
Produktbeschreibung	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[LK2H10 PHE5]

Molekulargewicht	68-75 kDa
Isotyp	IgG1
UniProt	P10645
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user