

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

**Melanoma Marker (MART-1 + Tyrosinase + gp100)(DT101+ BC199 + T311 + HMB45), CF647 conjugate, 0.1mg/mL, Clone: [DT101 BC199 T311 HMB45], Mouse, Monoclonal
BOT-BNC470704-100**

Artikelname	Melanoma Marker (MART-1 + Tyrosinase + gp100)(DT101+ BC199 + T311 + HMB45), CF647 conjugate, 0.1mg/mL, Clone: [DT101 BC199 T311 HMB45], Mouse, Monoclonal
Artikelnummer	BOT-BNC470704-100
Hersteller Artikelnummer	BNC470704-100
Alternativnummer	BOT-BNC470704-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant hMART-1 protein (DT101 & BC199), Extract of pigmented melanoma metastases from lymph nodes (HMB45), Recombinant tyrosinase protein (T311)
Konjugation	CF647
Produktbeschreibung	This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[DT101 BC199 T311 HMB45]
Molekulargewicht	20-22 kDa (doublet) (MART-1), 90-100 kDa (gp100), 70-80 kDa (Tyrosinase)
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user