

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **c-Myc(MYC909), CF647 conjugate, 0.1mg/mL, Clone: [MYC909], Mouse, Monoclonal BOT-BNC470909-500**

|                          |   |
|--------------------------|---|
| Artikelname              | c-Myc(MYC909), CF647 conjugate, 0.1mg/mL, Clone: [MYC909], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC470909-500   |
| Hersteller Artikelnummer | BNC470909-500   |
| Alternativnummer         | BOT-BNC470909-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human c-myc protein   |
| Konjugation              | CF647   |
| Produktbeschreibung      | This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [MYC909]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 62-64 kDa  |
| UniProt                | <a href="#">P01106</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |