

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NGFR (Nerve Growth Factor Receptor)(NTR/912), CF647 conjugate, 0.1mg/mL, Clone: [NTR/912], Mouse, Monoclonal BOT-BNC470912-100

| | |
|--------------------------|---|
| Artikelname | NGFR (Nerve Growth Factor Receptor)(NTR/912), CF647 conjugate, 0.1mg/mL, Clone: [NTR/912], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470912-100 |
| Hersteller Artikelnummer | BNC470912-100 |
| Alternativnummer | BOT-BNC470912-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Recombinant human p75 NGFR protein |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes a glycoprotein of 75 kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melano... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [NTR/912] |

| | |
|------------------------|---|
| Molekulargewicht | 75 kDa |
| UniProt | P08138 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |