

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **S100B(4C4.9 + S100B/1012), 0.2mg/mL, Clone: [4C4.9 S100B/1012], CF647, Mouse, Monoclonal BOT-BNC471204-500**

Artikelname	S100B(4C4.9 + S100B/1012), 0.2mg/mL, Clone: [4C4.9 S100B/1012], CF647, Mouse, Monoclonal
Artikelnummer	BOT-BNC471204-500
Hersteller Artikelnummer	BNC471204-500
Alternativnummer	BOT-BNC471204-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	Purified bovine brain S100 protein (4C4.9), Recombinant full-length human S100B protein (S100B/1012)
Konjugation	CF647
Produktbeschreibung	S100 belongs to the family of calcium binding proteins. S100A and S100B proteins are two members of the S100 family. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epi...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[4C4.9 S100B/1012]
Molekulargewicht	10-12 kDa
UniProt	<a href="#">P04271</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user