

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nucleolin(364-5 + NCL/902), CF647 conjugate, 0.1mg/mL, Clone: [364-5 NCL/902], Mouse, Monoclonal BOT-BNC471205-100

Artikelname	Nucleolin(364-5 + NCL/902), CF647 conjugate, 0.1mg/mL, Clone: [364-5 NCL/902], Mouse, Monoclonal
Artikelnummer	BOT-BNC471205-100
Hersteller Artikelnummer	BNC471205-100
Alternativnummer	BOT-BNC471205-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Lysate of SU-DHL-1 Nuclei (364-5), Recombinant human NCL protein (NCL/902)
Konjugation	CF647
Produktbeschreibung	Recognizes Nucleolin (NCL), the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleolar chromatin and pre-ribosomal particles. Human NCL...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[364-5 NCL/902]
Molekulargewicht	76 kDa (predicted), 100-110 kDa (apparent)
UniProt	P19338
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with mouse, rat or cow, others not known Immunohistology (formalin) Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user