

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD74(LN-2 + CLIP/813), CF647 conjugate, 0.1mg/mL, Clone: [LN-2 CLIP/813], Mouse, Monoclonal BOT-BNC471207-500

| | |
|--------------------------|---|
| Artikelname | CD74(LN-2 + CLIP/813), CF647 conjugate, 0.1mg/mL, Clone: [LN-2 CLIP/813], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471207-500 |
| Hersteller Artikelnummer | BNC471207-500 |
| Alternativnummer | BOT-BNC471207-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | SU-DHL-4 lymphoma cells (LN-2), Recombinant full-length human CD74 protein (CLIP/813) |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes a protein of ~35 kDa, identified as CD74. It is a type II transmembrane protein which binds to the peptide binding groove of newly synthesized MHC class II alpha/beta heterodimers and prevents their premature association with... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [LN-2 CLIP/813] |
| Molekulargewicht | 33-41 kDa |
| UniProt | P04423 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |