

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Desmocollin-2/3 (7G6), CF647 conjugate, 0.1mg/mL, Clone: [7G6], Mouse, Monoclonal BOT-BNC471430-100

| | |
|--------------------------|---|
| Artikelname | Desmocollin-2/3 (7G6), CF647 conjugate, 0.1mg/mL, Clone: [7G6], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471430-100 |
| Hersteller Artikelnummer | BNC471430-100 |
| Alternativnummer | BOT-BNC471430-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Human desmocollin-2 extracellular domain (exact sequence is proprietary) |
| Konjugation | CF647 |
| Produktbeschreibung | Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. The desmosome is subdivided into two regions. The plaque region lies adjacent to the plasma, and is believed to contain molecules ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [7G6] |
| Molekulargewicht | 110 kDa (DSC2), 100 kDa (DSC3) |
| UniProt | Q02487 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |