

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Catenin, beta (p120) (15B8), CF647 conjugate, 0.1mg/mL, Clone: [15B8], Mouse, Monoclonal BOT-BNC471443-500

| | |
|--------------------------|---|
| Artikelname | Catenin, beta (p120) (15B8), CF647 conjugate, 0.1mg/mL, Clone: [15B8], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471443-500 |
| Hersteller Artikelnummer | BNC471443-500 |
| Alternativnummer | BOT-BNC471443-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Bovine, Canine, Gallus, Human, Mouse, Rat |
| Immunogen | Recombinant chicken β -catenin |
| Konjugation | CF647 |
| Produktbeschreibung | Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [15B8] |

| | |
|------------------------|--|
| Molekulargewicht | 92 kDa |
| UniProt | P35222 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |