

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD15 / FUT4 (Reed-Sternberg Cell Marker) (FUT4/1478R), CF647 conjugate, 0.1mg/mL, Clone: [FUT4/1478R], Rabbit, Monoclonal BOT-BNC471478-500**

Artikelname	CD15 / FUT4 (Reed-Sternberg Cell Marker) (FUT4/1478R), CF647 conjugate, 0.1mg/mL, Clone: [FUT4/1478R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC471478-500
Hersteller Artikelnummer	BNC471478-500
Alternativnummer	BOT-BNC471478-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Purified human neutrophils
Konjugation	CF647
Produktbeschreibung	CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cell...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[FUT4/1478R]

Molekulargewicht	220 kDa
UniProt	<a href="#">P22083</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow cytometry 2-4 ug/million cells Immunofluorescence 2-4 ug/mL Immunohistology (formalin) 5-10 ug/mL Optimal dilution for a specific application should be determined by user