

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD99 / MIC2 (Ewing s Sarcoma Marker) (MIC2/1495R), CF647 conjugate, 0.1mg/mL, Clone: [MIC2/1495R], Rabbit, Monoclonal BOT-BNC471495-100**

Artikelname	CD99 / MIC2 (Ewing s Sarcoma Marker) (MIC2/1495R), CF647 conjugate, 0.1mg/mL, Clone: [MIC2/1495R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC471495-100
Hersteller Artikelnummer	BNC471495-100
Alternativnummer	BOT-BNC471495-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length MIC2 protein
Konjugation	CF647
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 27-32 kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which ar...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MIC2/1495R]

Molekulargewicht	27-32 kDa
UniProt	<a href="#">P14209</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 2-4 ug/mL Flow cytometry 2-4 ug/million cells Immunohistology (formalin) 5-10 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user