

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD8B (Cytotoxic / Suppressor T-Cell Marker)(CBB/1595), CF647 conjugate, 0.1mg/mL, Clone: [CBB/1595], Mouse, Monoclonal BOT-BNC471595-500

| | |
|--------------------------|---|
| Artikelname | CD8B (Cytotoxic / Suppressor T-Cell Marker)(CBB/1595), CF647 conjugate, 0.1mg/mL, Clone: [CBB/1595], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471595-500 |
| Hersteller Artikelnummer | BNC471595-500 |
| Alternativnummer | BOT-BNC471595-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Bovine, Goat, Sheep |
| Immunogen | Antibody was raised against bovine CD8b. |
| Konjugation | CF647 |
| Produktbeschreibung | The CD8 beta chain associates with the CD8 alpha chain to form the CD8 alpha/beta heterodimer expressed on the surface of a majority of thymocytes, and on peripheral cytotoxic alpha beta TCR T cells. This monoclonal antibody identifies cytotoxic/supp... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CBB/1595] |
| Molekulargewicht | 32 kDa |

| | |
|------------------------|--|
| UniProt | P10966 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |