

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Galectin-1 / Human Placental Lactogen (hPL) (GAL1/1831), CF647 conjugate, 0.1mg/mL, Clone: [GAL1/1831], Mouse, Monoclonal BOT-BNC471831-500

| | |
|--------------------------|---|
| Artikelname | Galectin-1 / Human Placental Lactogen (hPL) (GAL1/1831), CF647 conjugate, 0.1mg/mL, Clone: [GAL1/1831], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471831-500 |
| Hersteller Artikelnummer | BNC471831-500 |
| Alternativnummer | BOT-BNC471831-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Galectin-1 protein fragment (aa12-108) (exact sequence is proprietary) |
| Konjugation | CF647 |
| Produktbeschreibung | Galectin-1 is a member of the beta-galactoside-binding family and is a dimeric protein of 14 kD participating in a variety of normal and pathological processes, including cancer progression. Galectin-1 can affect the proliferation of normal and malig... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [GAL1/1831] |
| Molekulargewicht | 14 kDa |
| UniProt | P09382 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL ELISA: 2-4 ug/mL for coating order Ab without BSA Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |