

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/294), CF647 conjugate, 0.1mg/mL, Clone: [rNX2/294], Mouse, Monoclonal BOT-BNC471836-100

| | |
|--------------------------|---|
| Artikelname | NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/294), CF647 conjugate, 0.1mg/mL, Clone: [rNX2/294], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471836-100 |
| Hersteller Artikelnummer | BNC471836-100 |
| Alternativnummer | BOT-BNC471836-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Gallus, Human, Mouse, Rat |
| Immunogen | Human full-length recombinant NKX2.2 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [rNX2/294] |

| | |
|------------------------|---|
| Molekulargewicht | 40-50 kDa |
| UniProt | O95096 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |